

ON THE MAGNETIC FIELDS AND PARTICLE ACCELERATION IN CAS A

JACCO VINK¹

Columbia Astrophysics Laboratory, Columbia University, MC 5247, 550 W 120th street, New York, NY 10027, USA

jvink@astro.columbia.edu

J. MARTIN LAMING

Naval Research Laboratory, Code 7674L, Washington DC 20375, USA

jlaming@ssd5.nrl.navy.mil

Draft version February 2, 2008

ABSTRACT

We investigate the non-thermal X-ray emission from Cas A, using *BeppoSAX*, *CGRO-OSSE*, and *Chandra* data. For the hard X-ray continuum we test the model proposed by Laming, which invokes non-thermal bremsstrahlung from electrons accelerated by lower hybrid plasma waves. The justification for this model comes from our determination of a lower limit to the average magnetic field of $B > 0.5$ mG. For such high magnetic fields the synchrotron losses are severe enough that most of the electron populations responsible for the radio emission have maximum electron energies well below the limit for which X-ray synchrotron emission is important. However, we do suggest that the rim surrounding Cas A, seen in *Chandra* continuum images, is X-ray synchrotron emission. The width of this rim of $1.5''$ to $4''$, can be used to infer the magnetic field near the shock front, for which we estimate $B = 0.08 - 0.16$ mG, and electron energies of $\sim 57 - 40$ TeV. This magnetic field strength is lower than the average magnetic field, but higher than what may be expected from shocked interstellar medium, suggesting either a high magnetic field in the wind of the progenitor, or rapid, post shock, field amplification by non-linear growth of plasma waves. Combining the two magnetic field measurements we have constructed a simple two zone model. Most of the radio emission comes from inside Cas A where the magnetic field is strong. In contrast, the inverse Compton emission is dominated by emission from near the shock front. Only for our lower limit on the magnetic field strength near the shock front is it possible to explain the recent detection of TeV emission by *HEGRA* with inverse Compton, for which, in addition, we have to assume a rather high far infrared photon density that should be twice as high as our best estimate of ~ 70 cm⁻³. Pion decay is therefore likely to be the dominant emission from Cas A at TeV energies.

Subject headings: shock waves – magnetic fields – gamma rays: observations – X-rays: observations
 individual (Cassiopeia A) – supernova remnants –

1. INTRODUCTION

Supernova remnants have long been thought to be the source of cosmic rays with energies up to at least 10^{15} eV, where there is a break in the observed cosmic ray spectrum, usually referred to as the “knee”. The recent discoveries of X-ray synchrotron and TeV emission from some supernova remnants support this long held belief by confirming that electrons are accelerated to energies of at least a few TeV (Reynolds & Keohane 1999). This is still somewhat short of the “knee”, but unlike for ions, the maximum electron energy may be limited by radiation losses.

SN 1006 has become the canonical example of a supernova remnant with X-ray emission dominated by synchrotron radiation from shock accelerated electrons (Koyama et al. 1995). The presence of electrons in excess of 10 TeV is further supported by the detection of TeV emission with the *CANGEROO* Čerenkov detector (Tanimori et al. 1998). The observed TeV emission is most likely inverse Compton scattering by the same population of electrons that is responsible for the X-ray continuum (Tanimori et al. 1998), although there is some debate about this interpretation (Berezhko et al. 2002).

Another often mentioned candidate for an X-ray syn-

chrotron emitting supernova remnant is Cas A. Its soft X-ray emission is not continuum dominated, but the observed emission above 20 keV cannot be explained by thermal bremsstrahlung, and has therefore been attributed to synchrotron radiation (The et al. 1996; Allen et al. 1997; Favata et al. 1997). The detection of TeV emission by *HEGRA* is consistent with this interpretation, but pion decay associated with relativistic ions is an equally valid interpretation (Aharonian et al. 2001).

For Cas A the interpretation of both the hard X-ray and the TeV emission is critically dependent on the magnetic field strength. Equipartition arguments to explain the radio luminosity imply magnetic fields of $B \sim 0.4 - 2$ mG (Longair 1994; Wright et al. 1999), far in excess of the canonical value for the interstellar medium of $3 \mu\text{G}$. The typical loss time of an electron with energy E , due to synchrotron radiation is (e.g. Reynolds 1998):

$$\tau_{\text{loss}} = \frac{635}{B^2 E} \text{ s.} \quad (1)$$

For a magnetic field of 1 mG, a particle with $E = 8$ TeV loses its energy in 2 yr. For Cas A, 8 TeV is the maximum energy needed to explain the hard X-ray emission (Reynolds & Keohane 1999). The two year lifetime is much shorter than the lifetime of Cas A, which is probably the

¹ Chandra fellow

remnant of a supernova that exploded in AD 1680 (Ashworth 1980). Similarly a high magnetic field lowers the number density of relativistic electrons for a given radio brightness, which in turn implies a low inverse Compton contribution to the observed TeV emission.

One way out of this situation would be to assume that electrons are continuously being reaccelerated, but that seems unlikely. The radio morphology of Cas A shows a bright ring of radius $\sim 1.8'$, whereas the shock front, the primary site for cosmic ray acceleration, is at a radius of $\sim 2.6'$. Moreover, detailed radio spectral index studies do not find evidence of reacceleration in bright radio knots, which are probably ejecta fragment penetrating the shell (Anderson & Rudnick 1996; Wright et al. 1999). These studies rather suggest that brightness variations are the result of variations in the magnetic field, which light up the diffuse relativistic electron population in various degrees.

Here we test an alternative interpretation of the hard X-ray emission with *BeppoSAX*-PDS data, namely bremsstrahlung from electrons accelerated by lower hybrid waves, which was suggested by Laming (2001a,b). An additional argument for testing such a model is that *XMM-Newton* observations show that even above 10 keV a substantial fraction of the continuum emission is not associated with the forward shock (Bleeker et al. 2001). The *BeppoSAX* data, in combination with *CGRO*-OSSE data, also provides an upper limit on the bremsstrahlung from the electron cosmic ray injection spectrum, from which a new lower limit on the average magnetic field strength can be obtained.

Nevertheless, we argue that X-ray synchrotron radiation does exist in Cas A, and is coming from close to the forward shock, where it shows up as a narrow rim surrounding the remnant, observed in *Chandra* X-ray continuum images (Gotthelf et al. 2001). Interpreting the width of the rim as a typical length scale for synchrotron losses, we are able to constrain the magnetic field near the shock front. Our analysis supports the idea that the average magnetic field is close to its equipartition value, whereas we find that near the shock front the magnetic field is lower, but still an order of magnitude higher than what may be expected from shocked interstellar medium ($\sim 10^{-5}$ G). We will discuss the implications of these values for the hard X-ray and TeV emission.

2. DATA AND OBSERVATIONS

Our analysis of the X-ray emission of Cas A is based on *BeppoSAX*-PDS data and archival *CGRO*-OSSE and *Chandra* data. The *BeppoSAX*-PDS (Frontera et al. 1997) and *CGRO*-OSSE (Johnson et al. 1993) instruments both consist of four phoswich detectors (NaI(Tl)/CsI(Na) scintillation detectors), behind two rocking collimators. They differ, however, substantially in energy range, field of view, and effective area. The *BeppoSAX*-PDS instruments cover an energy range of 15–300 keV, whereas this is 50 keV to 10 MeV for *CGRO*-OSSE.

BeppoSAX observed Cas A for 500 ks in May and June 2001 in order to observe the 67.9 keV and 78.4 keV nuclear decay lines of ^{44}Sc , the nuclear decay product of ^{44}Ti . The detection of these lines at the $3\text{--}5\sigma$ level, depending on the

hard X-ray continuum model, was reported by Vink et al. (2001), which provides more details about the data set. Due to the rocking collimators the effective exposure time is about half the observation time. Together with archival *BeppoSAX* data the spectra used for our analysis here is based on a total effective PDS exposure of 311 ks.

The spectrum, as presented in Vink et al. (2001), clearly shows the hard X-ray spectrum with a bump associated with the ^{44}Sc line emission. Marginal (2σ) residual emission around 60 keV is possibly caused by fluorescent tantalum K shell emission, as a result of cosmic rays interacting with the collimator, which consists of tantalum with a thin copper-tin bi-layer. Some residual emission above 100 keV is not easily explained (cf. Fig. 1 & 2). It may be intrinsic to Cas A, but some instrumental contamination seems more likely. The residual emission peaks around 200 keV, but is not narrow enough to be caused by fluorescent line emission from any particular material. There are no other similar deep *BeppoSAX* observations with a moderate hard X-ray brightness like Cas A, which would allow us to identify possible instrumental features at this level.

Although the residual emission above 100 keV is statistically not very significant, one of our concerns is to obtain an upper limit on the emission above 100 keV in order to obtain a lower limit to the average magnetic field in Cas A. We therefore included archival *CGRO*-OSSE spectra in our analysis. Moreover, the effective area of the instruments peaks around 250 keV, making it useful for constraining the emission of Cas A above 100 keV. Cas A was observed several time by *CGRO*-OSSE (see The et al. 1996). We added all available Cas A spectra and averaged the corresponding response matrices. The observations were made during the viewing periods 34 – 815 with a total exposure of 909 ks.² The total exposure time was 909 ks.

In addition to hard X-ray data, we use a deep (50 ks) archival *Chandra* observation with the ACIS-S3 chip to complement our analysis. The observation was made in January 2000. The same data were used by Hwang et al. (2000). For our analysis we used level 2 archival data products, generating spectra and calibration products with the standard *Chandra* software package *Ciao* v1.1.0.

3. THE LOWER HYBRID ELECTRON ACCELERATION MODEL

Cas A's morphology indicates a complicated, turbulent hydrodynamical structure. Radio, optical and X-ray images all show many bright knots on top of more diffuse emission. As the forward or reverse shocks strike and illuminate these features, secondary shocks will split off from these primary shocks and propagate back into the already shocked circumstellar or ejecta plasma. In this way shocks may propagate across the likely region of high magnetic field at the contact discontinuity, where field amplification by Rayleigh-Taylor instability may occur. The acceleration of electrons by lower hybrid waves generated by these shocks has been considered by Laming (2001a,b). Bremsstrahlung emitted by these accelerated electrons was shown to provide a very good match to the hard X-ray continuum of Cas A, using the ~ 50 ksec of *BeppoSAX*-PDS data that were available at the time.

² In addition to the observations listed in The et al. (1996), this means VP 617, 715, 810, 814, and 815.

Lower hybrid waves (LHW) are electrostatic ion oscillations directed within an angular range $\pm\omega_{pi}/\omega_{pe}$ of the direction perpendicular to the magnetic field, where ω_{pe} and ω_{pi} are the electron and ion plasma frequencies respectively. The electron screening that would usually damp such oscillations is inhibited by the magnetic field, giving a criterion on the electron temperature that the electron gyroradius be smaller than the wavelength divided by 2π . Since $\omega/k_{\perp} \ll \omega/k_{\parallel}$ the wave can simultaneously be in resonance with ions moving across the magnetic field lines and electrons moving along them. Consequently energy equilibration between ions and electrons may proceed at a much faster rate than by Coulomb collisions alone, and saturate when the electron thermal pressure is equal to the magnetic pressure. Such processes have been considered in the context of two-temperature accretion flows by Begelman & Chiueh (1988). Here cross field ion motions are produced by a combination of curvature and diamagnetic drift, and the magnetic field is presumably generated by a magneto-rotational instability (Balbus & Hawley 1991, 1998; Quataert et al. 2002), the critical issue being whether sufficient electron-ion equilibration can occur to destroy ADAF or similar solutions for the accretion flow. In dilute plasmas where the electron acceleration time is much shorter than the electron-electron collision time, a non-Maxwellian electron distribution in the component of velocity along the magnetic field can result. This has been modeled numerically using particle-in-cell simulations by Shapiro et al. (1999) for application to cometary X-rays, and analytically by Vaisberg et al. (1983) and Krasnosel'skikh et al. (1985). Lower hybrid waves generated by a modified two-stream instability and the resulting accelerated electrons were all observed at Comet Halley (Klimov et al. 1986; Gringauz 1985). Bingham et al. (2000) considered LHW in supernova remnants and Laming (2001a,b) applied the analytic formulations, which agree well with the numerical work, to the case of accelerated electrons in Cas A. The non-Maxwellian electron distribution function depends on two parameters, the overall normalization (i.e. the fraction of plasma electrons that are accelerated) and the maximum electron energy in the distribution. In our fitting we treat these both as free parameters. A demonstration of how these variables are related to other physics of the Cas A plasma, and hence could be determined is given in the Appendix.

We fitted the predicted spectrum, calculated as in Laming (2001a), but with a relativistic bremsstrahlung cross section taken from Nozawa, Itoh, & Kohyama (1998), to the observed *BeppoSAX*-PDS X-ray spectrum. Apart from the non-thermal bremsstrahlung spectrum several other components, which are likely to contribute to the hard X-ray emission, are included in the spectral model. At the low end of the hard X-ray spectrum, the tail of the thermal bremsstrahlung with $T_e = 4 \times 10^7$ K is expected to contribute to the spectrum. In addition, ^{44}Sc line emission at 68 and 78 keV is expected, as a result of the decay of ^{44}Ti synthesized during the explosion (Vink et al. 2001; Iyudin et al. 1994). The line fluxes of the two nuclear transitions are expected to be almost identical. A broadening corresponding with a velocity dispersion of 5000 km s^{-1} was included. A residual feature around 60 keV in the ob-

served spectrum may be attributed to cosmic ray induced fluorescence from the tantalum collimator, and was modeled by delta functions corresponding to Ta $K\alpha$ (57 keV) and $K\beta$ (65 keV) with a ratio of 4:1.³ The spectrum and best fit model are shown in Fig. 1 and model parameters are given in Table A1.




FIG. 1.— *BeppoSAX*-PDS spectrum of Cas A with the best fit ^{44}Sc /non-thermal bremsstrahlung model. The individual emission components are: ^{44}Sc line emission (red), non-thermal continuum (blue), thermal continuum (green) and possible line contamination from collimator material (tantalum, cyan). The observed count rate in each channel has been divided by the effective area in order to yield approximate flux densities.

The parameter values indicate that electrons are possibly accelerated up to ~ 95 keV. The model gives a very good fit to the spectrum up to 100 keV. However, it does not describe the emission observed above 100 keV. As discussed above, the data above 100 keV may be less reliable, and the low flux point around 100 keV suggest that the actual continuum emission is indeed curved, as predicted by the LHW model.

In Table A1 we also list for comparison the parameters for a simple power law continuum spectrum, and the X-ray synchrotron models available in the spectral fitting package *xspec* (Arnaud 1996), *srcut* and *sresc* (Reynolds 1998; Reynolds & Keohane 1999). The *srcut* model is a very simplified model of X-ray synchrotron emission from a relativistic electron distribution with an exponential cut-off. In reality the cut-off energy is likely to vary from place to place, giving a more gradual decrease in slope. Moreover, as explained in the introduction, for the bulk of the relativistic electrons, at a remote distance from the shock front, the maximum electron energy is probably far below what is needed to produce X-ray synchrotron emission. The *sresc* model takes these spatial effects into account, assuming a Sedov model for the evolution of the supernova remnant. Although, *sresc* is more physical, it may not be readily applicable to Cas A, which has large magnetic field gradients and is not well described by the standard Sedov model (Vink et al. 1998).

The LHW model is in a similar way somewhat oversimplified, as also here it is likely that maximum energies vary within the remnant, depending on local conditions such as the local shock velocity, obliquity and ion composition. Nevertheless, given the limited number of parameters, the model fits very well. Note that the spectral slope between

³ See the X-ray data booklet (Thompson et al. 2001).

20-50 keV, is a property of the model, and not a free parameter.

Finally, note that using the LHW model results in 50% higher ^{44}Sc line fluxes, than using a power law continuum, a result of the continuum curvature predicted by both models. The higher ^{44}Sc line flux is close to the latest *CGRO-COMPTEL* measurements of the ^{44}Ca line flux (Schönfelder et al. 2000), and translates into a synthesized ^{44}Ti mass of $1.8 \times 10^{-4} M_{\odot}$, for a distance of 3.4 kpc (Reed et al. 1995) and a ^{44}Ti decay time of 85.4 yr (Ahmad et al. 1998; Görres et al. 1998; Norman et al. 1998).

4. A NEW LOWER LIMIT TO THE AVERAGE MAGNETIC FIELD

The LHW model is attractive to consider for Cas A because of its high magnetic field strength, which is estimated to be ~ 1 mG, based on the assumption of equipartition of the relativistic particle population and the magnetic field energy (e.g. Longair 1994). Although there is no compelling physical reason why the equipartition assumption should be valid, estimates based on the absence of gamma-ray emission from Cas A supports a value of the magnetic field in excess of what might be expected from a shock compression of the interstellar magnetic field (Cowsik & Sarkar 1980). The most recent estimate of the magnetic field is based on the upper limit obtained by *CGRO-EGRET*, suggesting $B > 3.5 \times 10^{-4}$ G (Esposito et al. 1996; Atoyan et al. 2000). This value is derived by comparing the radio synchrotron flux density with the expected bremsstrahlung from the relativistic electrons responsible for the radio emission.

Here we use essentially the same method, but less directly, as we use an extrapolation of the relativistic electron population to energies observed by *BeppoSAX*-PDS and *CGRO-OSSE*. The extrapolation used is similar to the one by Asvarov et al. (1990), and assumes that the electron momentum spectrum is a power law distribution, as is expected for an electron population generated by first order Fermi acceleration (Bell 1978). This means that the energy spectrum flattens at non-relativistic energies. The electron cosmic ray injection spectrum is not well known, but models indicate that at lower energies the spectrum is steeper than expected from extrapolating the relativistic spectrum (Bykov & Uvarov 1999). In that case our upper limit on the cosmic ray normalization is too high, which translates into too low a lower limit.

Of more concern is the role of energy losses on the shape of the spectrum, as it will tend to flatten the low energy end of the spectrum. We checked the signature of adiabatic losses on the spectrum, and found that it does affect the spectrum mostly at mildly relativistic energies, where it results in a steeper spectrum at low energies, is flatter at intermediate energies and then assumes the original power law index at extreme relativistic energies. We did not take the steepening into account, as we do not know how much the average electron spectrum is affected by it. But if there is a steepening, it makes our estimates too conservative. The most dominant other loss factor between 100 and 1000 keV in the ionized plasma of the remnant is electron-electron collisions. For an electron temperature of 4×10^7 K and density of 20 cm^{-3} , the loss time of a 100 keV electron is ~ 200 yr, rapidly increasing to ~ 500

yr for 200 keV (Huba 1994). These loss times are comparable to or larger than the average plasma age of Cas A, indicating that non-adiabatic losses are likely to be small.

The synchrotron flux density for a relativistic electron power law distribution, with energy index q , is given by (Blumenthal & Gould 1970):

$$F_{\nu} = 1.70 \times 10^{-21} a(q) \frac{\kappa V}{4\pi d^2} B^{\frac{q+1}{2}} \left(\frac{6.26 \cdot 10^{18}}{\nu} \right)^{\frac{q-1}{2}} \quad (2)$$

where κ is the particle normalization, V the emitting volume, d the distance, B the magnetic field, and $a(q)$ is a factor tabulated in Blumenthal & Gould (1970).

Bremsstrahlung from the same population of electrons, which flattens at non-relativistic energies, scales with the factor $\Sigma_i n_i Z_i^2 \kappa V / 4\pi d^2$, where $\Sigma_i n_i Z_i^2$ indicates the sum of densities of all the ion species with charge Z_i . So from the combination of synchrotron emission and bremsstrahlung, and an estimate of $\Sigma_i n_i Z_i^2$, we can estimate κ and B .

The radio spectrum of Cas A has a spectral index of $\alpha = -0.78$ and based on a compilation of radio data (e.g. Baars et al. 1977) we estimate a radio flux density of 2522 Jy at 1 GHz for the epoch 2000. The spectral index indicates an electron energy spectrum with $q = 2.56$ at relativistic energies and flattening to 1.77 at non-relativistic energies, resulting in an X-ray photon index of -2.2. This is at odds with the observed spectral index of -3.3, which means that 100 keV must have a different origin, taken here to be bremsstrahlung from LHW accelerated electrons. We adopt the LHW model, and try to obtain an upper limit on the emission from an additional component with a power law slope of -2.2 from the combined *BeppoSAX* and *CGRO-OSSE* data (Fig. 2).




FIG. 2.— Best fit model that combines the LHW/ ^{44}Sc with the electron cosmic ray injection spectrum. Shown are the *BeppoSAX*-PDS (black) and *CGRO-OSSE* (red) spectrum of Cas A.

The best fit normalization for such a power law spectrum is $(4.0 \pm 1.5) \times 10^{-7} \text{ ph s}^{-1} \text{ keV}^{-1} \text{ cm}^{-2}$ at 100 keV, with a 2σ upper limit of $6.2 \times 10^{-7} \text{ ph s}^{-1} \text{ keV}^{-1} \text{ cm}^{-2}$. We measured the electron spectrum normalization more directly by fitting a model of bremsstrahlung from a relativistically correct power law momentum spectrum, using the gaunt factors proposed by Haug (1997). The upper limit on the bremsstrahlung normalization obtained this way is $\kappa \Sigma_i n_i Z_i^2 V / 4\pi d^2 < 65$. Soft X-ray observations imply $\Sigma_i n_i Z_i^2 \simeq 20$ (Vink et al. 1996; Willingale et al. 2001b), which, combined with the bremsstrahlung normalization,

gives for the average magnetic field $B > 0.5$ mG.⁴ This is higher than the value derived by Atoyan et al. (2000) from the *CGRO-EGRET* upper limit, especially taking into account that they assumed a higher value for $\Sigma_i n_i Z_i^2$. Note that the above lower limit is not a volume average, but based on the ratio $\langle \kappa B^{(q+1)/2} \rangle / \langle \kappa \Sigma_i n_i Z_i^2 \rangle$. The derived lower limit is in good agreement with equipartition of magnetic and cosmic ray energy.

5. THE MAGNETIC FIELD NEAR THE SHOCK FRONT

The high average magnetic field in Cas A is inconsistent with simple shock compression of the interstellar magnetic field, and indicates that the magnetic field inside the remnant must have been enhanced by turbulence. A likely place for turbulence is the contact discontinuity, which may explain why the radio emission has a bright ring of emission, as this is the region in the remnant where magnetic field amplification is important. A similar argument was used by Anderson & Rudnick (1996) to explain why radio knots have the same radio spectral index as the surrounding diffuse emission; the enhanced magnetic field lights up the background electron population. One may wonder then, what the initial interstellar magnetic field was, before turbulent magnetic field amplification.

An estimate can be derived from the observation of the shock front of Cas A by *Chandra*. In the X-ray continuum band from roughly 4.5-6 keV Cas A seems bound by a narrow rim of emission, which marks the onset of radio emission. The narrowness of the rim indicates strong limb brightening of a layer, which much have some special emission properties. The fact that the radio and X-ray morphology are dissimilar as far as this rim is concerned, seems to argue against a synchrotron origin for the X-ray emission, see e.g. Gotthelf et al. (2001). But this is not true, if the width of the rim is a result of the limited lifetime of the ultra-relativistic electrons, as soon as they are swept away from the shock front. Further inside the shock the maximum energy of the electrons has so much decreased that no X-ray synchrotron emission is produced anymore. This is not the case for the radio emission, caused by electrons with much lower energies, for which synchrotron and inverse Compton losses are negligible (e.g. Wright et al. 1999). This behavior can be seen in the models by Reynolds (1998), in which the synchrotron emitting shell is thinner at higher frequencies. The steady increase in radio emission toward the bright ring of emission may be the effect of accumulation of relativistic electrons swept away from the shock front, and a strong gradient of the magnetic field.

In this interpretation most of the electron populations, which reside outside the X-ray rims, have a maximum energy cut-off well below 1 TeV, unless reacceleration is important. However, radio spectral index measurements do not indicate any reacceleration of electrons by ejecta knots, which would likely lead to different indices of radio knots and diffuse emission (Anderson & Rudnick 1996). Note that this argues also against the use for Cas A of the simple maximum curved synchrotron model employed by Reynolds & Keohane (1999), which merely assumes that all electron populations have the same energy cut-off, and

the magnetic field is constant.




FIG. 4.— *Chandra* ACIS-S3 X-ray spectra of the narrow rim of emission (lower spectrum) and from emission just inside the rim ($\times 10$). Note the lower equivalent width of the line emission from the spectrum of the rim (cf. Table A2).

The rim is much narrower than the $\sim 1/12$ th shock radius expected for plasma shocked with a factor 4 compression ratio. That they are indeed consistent with X-ray synchrotron emission can be exemplified by a comparison of the X-ray continuum flux and radio flux of part of the rim in the northeast (see Fig. 3, and Table A2). We used archival *Chandra* data to obtain a spectrum from this filament and determined that the X-ray continuum flux between 4 and 6 keV is 1.0×10^{-12} erg s $^{-1}$ cm $^{-2}$ (2.1×10^{-7} Jy at 1.2×10^{18} Hz). We normalized an archival VLA radio map (epoch 1987) to a flux of 2522 Jy and extracted a flux density from the same region. As the shock front may have moved between 1987 and 2001 by as much as 5'', we extracted also a flux from the same region, but moved 5'' toward the center of the remnant. This yielded a flux density estimate between 3 Jy and 6 Jy at 1 GHz, corresponding to a broad band spectral index of $\alpha = -1.15$ (the factor two uncertainty in flux density is irrelevant over a frequency range of 9 decades). This is steeper than the radio spectral index of 0.78 and thus indicates that the X-ray flux density is below the extrapolation of the radio spectrum; consistent with the interpretation that part of the X-ray emission is synchrotron emission.

Indeed, not all X-ray emission is synchrotron, as the narrow rim displays line emission from Mg, Si and S, which should be accompanied by bremsstrahlung as well. The line emission from the rim is, however, weaker and the spectrum harder, than for the spectrum of a region just inside of the rim (Table A2). For example fitting a thermal non equilibrium model with an additional power law gives a spectral index of -2.2, whereas away from the rim the spectral index is unconstrained, but steeper than -4. Moreover, it is hard to find another interpretation for the rim than synchrotron emission, as only energy losses can explain the narrow features. For a hot plasma the losses are small and the gradual ionization and possible equilibration of electrons and ions behind the shock would lead to a steady increase in intensity behind the shock, not in a sudden onset of emission, followed by a rapid decline.

⁴ This replaces the value reported in Vink & Laming (2002), in which we missed a factor 4π in the synchrotron normalization (eq. 2, c.f. Ginzburg & Syrovatski 1965).




FIG. 3.— Detail of a VLA radio image (left) and *Chandra*-ACIS X-ray image from a compilation of continuum dominated bands between 2.7 - 9 keV. The spectral extraction regions are indicated in the *Chandra* image (NE rim and NE inside rim).

Interestingly, identifying the rims with X-ray synchrotron emission from recently accelerated electrons makes this aspect of Cas A very similar to SN 1006, where the X-ray synchrotron emission is coming from a similarly narrow rim of emission, as shown by archival *Chandra* data. For SN 1006, however, the continuum emission from the rims are dominating the total continuum emission, whereas for Cas A the X-ray continuum morphology shows that a substantial fraction of the X-ray continuum is associated with the bright ejecta shell. In SN 1006 the synchrotron emission comes mainly from the northeastern and southwestern limb, which is usually interpreted as caused by the large scale orientation of the magnetic field (Reynolds 1998). For Cas A, however, we do not see such axi-symmetry. This implies a much more randomly oriented magnetic field in the surrounding medium.

Assuming that the limb brightened rim emits indeed X-ray synchrotron radiation, we can derive from the typical width of the rim the magnetic field behind the shock front and the typical electron energy responsible for the synchrotron radiation. The electron energy may be associated with the typical exponential cut-off energy of the electron spectrum, but the very nature of the losses make it likely that a range of cut-off values exist. In an equilibrium situation synchrotron and inverse Compton losses will steepen the spectrum by a $\Delta q = -1$ (Longair 1994), but as soon as the electrons are sufficiently far removed from the shock front the spectrum is more likely to have an exponential cutoff. An exponential cut off is also more likely, if the spectrum near the shock front is age limited instead of loss limited (Reynolds 1998).

The rims have a typical width of $1.5''$ - $4''$ and clearly stand out above the background and diffuse emission

(Fig. 5). We can estimate a typical loss time from the widths by combining them with the plasma velocity away from the shock front. The shock velocity in Cas A has been estimated from proper motion studies using Einstein and ROSAT HRI X-ray images, and is thought to be $v_s = 5200 \text{ km s}^{-1}$ (Vink et al. 1998). The velocity of plasma away from the shock front, for a standard shock compression ratio of 4, is $0.25v_s = 1300 \text{ km s}^{-1}$. This value is consistent with X-ray Doppler shifts, which imply a velocity relative to the shock front of $\Delta v \sim 1400 \text{ km/s}$ (Willingale et al. 2001a). For a distance of 3.4 kpc the measured widths of the rims correspond to a typical loss time of 18 to 50 yr.

The energy history of an electron with initial energy E_0 , suffering synchrotron losses, is:

$$E(t) = \frac{E_0}{1 + E_0 B^2 t / 635}. \quad (3)$$

After a time τ (eq.1), the electron will have an energy $\frac{1}{2}E_0$, whereas higher energy electrons, after the same amount of time, will have energies $E'(\tau) \leq E_0$. The relation between the peak emitting frequency and electron energy is given by (Ginzburg & Syrovatski 1965):

$$\nu_{peak} = 1.8 \times 10^{18} B_{\perp} E^2. \quad (4)$$

In reality we have to take into account a range of frequencies, as the the synchrotron emission function is broad.

The rim is best visible in the X-ray continuum dominated band of 4-6 keV, and in some narrow bands in between line emission features. Taking 5 keV ($1.2 \times 10^{18} \text{ Hz}$) as the typical photon energy, we can combine eq. 4 with eq. 1 to obtain a typical electron energy of $E = 65 \text{ erg}$ to 91 erg and $B = 0.16 \text{ mG}$ to 0.08 mG . The power slope of the Chandra spectrum near the shock is -2.2 , which indicates that the photons are emitted by electrons close

to the break in the electron distribution. In fact using the *srcut*-model (cf. §3), we derive for $B = 0.1$ mG, $E_{break} \sim 25$ erg. However, the spectrum is likely to be an integration of spectra with different break energies. So close to the shock front E_{break} is likely to be higher.




FIG. 5.— Emission profiles of the rim in the southeast (left) and northeast. The profiles were extracted using strips of 20 pixels wide ($9.8''$) from a *Chandra* image, which was a combination of continuum dominated bands between 2.7 and 9 keV.

In Fig. 6 we have illustrated our estimates of magnetic field strength and electron energy graphically. Apart from synchrotron losses we also have included inverse Compton losses. For this we have estimated the photon energy density from the cosmic microwave background and the far infrared emission from Cas A itself, the latter being the most dominant factor. From the *IRAS* infrared fluxes listed by Braun (1987) we estimate a photon energy density of $u = 4.7 \times 10^{-12}$ erg cm $^{-3}$, corresponding to an equivalent magnetic field of $B = 1.1 \times 10^{-4}$ G.

The estimates of the magnetic field near the shock front give values of roughly a fifth of the average magnetic field strength and indicate a preshock magnetic field of $\sim 2 - 4 \times 10^{-5}$, a factor ten higher than the canonical interstellar medium value. This may either hint at a higher magnetic field surrounding Cas A, for instance because the wind of the Wolf-Rayet progenitor carried a high magnetic field, or the presence of accelerated particles near the shock front has given rise to magnetic field amplification due to non-linear plasma wave growth. The first hypothesis was suggested by Biermann & Cassinelli (1993) in order to explain the acceleration of cosmic ray by supernova remnants up to 10^{16} eV for which high magnetic fields are necessary. The second hypothesis was suggested by Lucek & Bell (2000) for identical reasons.




FIG. 6.— The shaded area indicates the most likely values of E_{max} and B just behind the shock front. This area is formed by the two dashed lines, which mark the electron energies that contribute to the continuum emission around 5 keV, allowing for emission at 1/3rd and 10 times the peak frequency for synchrotron emission, and the solid lines, which are formed by assuming the width of the rims is determined by radiative losses (decay times of 18 to 50 yr and a shock velocity of 5200 km/s are assumed). The width of the rims should be at least be larger than twice the gyro-radius. This means that for a typical width of $3''$, the hatched region is excluded.

6. THE MAGNETIC FIELD STRENGTH AND THE NATURE OF THE TEV EMISSION

The measured magnetic field strengths have important consequences for the gamma ray emission from Cas A. For instance, our lower limit on the average magnetic field strength implies that the gamma ray emission from bremsstrahlung is almost an order of magnitude below the *CGRO-EGRET* upper limit.

It is also understood that for the high magnetic field in Cas A, it is unlikely that the detected TeV emission is inverse Compton emission (Aharonian et al. 2001; Atoyan et al. 2000). We have investigated this important issue by constructing a two zone model. Zone 1 is responsible for the bulk of the radio emission, such as the bright radio ring. It should have $B > 0.5$ mG according to lower limit on the magnetic field. As the electron population in this zone is old, no X-ray synchrotron emission is produced. We adopt a somewhat arbitrary maximum electron energy of 0.1 TeV, but in reality a range of values exist inside the remnant. The inverse Compton emission from this zone is negligible.

Zone 2 is responsible for the X-ray synchrotron emission from the rim and has $B \sim 0.08 - 0.16$ mG and $E_{max} \sim 40 - 60$ TeV. Based on our flux measurements, and assuming spherical symmetry, we estimate that the X-ray synchrotron emission should be less than one third of the X-ray emission around 5 keV, but as we focused on a particular bright part of the rim, 1/10th to 1/20th seem more likely values. Moreover, for $B = 0.08$ mG and an X-ray synchrotron flux contribution of 1/3rd, the non-thermal bremsstrahlung flux at 100 keV is above the measured upper limit. This criterion is relaxed for higher magnetic

fields near the shock front.




FIG. 7.— A simplified two zone model of the broad band radiation from the cosmic ray electron population, i.e. synchrotron radiation (dotted line), bremsstrahlung (dashed-dotted), and inverse Compton emission (dashed line). The solid line gives the total emission from both zones. The parameters near the shock front (zone 2) were chosen to be $B = 1.0 \times 10^{-4}$ G, $E_{max} = 55$ TeV. $B = 0.5$ mG for zone 1. The data points give the flux at 1 GHz, measurements by ?, X-ray continuum emission (*ASCA* and *BeppoSAX*), *CGRO-COMPTEL* (Strong et al. 2000), the *CGRO-EGRET* upper limit (Esposito et al. 1996), and *HEGRA* (Aharonian et al. 2001).

This two zone model is different from the two zone model proposed by Atoyan et al. (2000), for which the two zones relate to the radio knots and diffuse radio emission. We adopt here the view of Anderson & Rudnick (1996), that there are no intrinsic differences between the electron cosmic ray populations of the radio knots and the diffuse emission, except that in the radio knots the magnetic field is stronger. Note that the electron cosmic ray density is higher in zone 2. Zone 1 is brighter at radio wavelength due to the stronger magnetic field.

In order to estimate the inverse Compton emission we need to estimate the photon density caused by Cas A's far infrared emission. *IRAS* flux measurements (Braun 1987) indicate a total photon flux of $F_{ph} = 3.9 \times 10^5$ ph s $^{-1}$ cm $^{-2}$, centered around 10^{13} Hz. The far infrared emission is associated with ejecta (Lagage et al. 1996). Approximating the geometry by a sphere with radius similar to the ejecta, $r = 1.8$ pc, and assuming that half the number of photons spend an average time of r/c inside the sphere, we obtain for the photon density⁵:

$$n_{ph} = \frac{3}{2} \frac{d^2}{r^2} \frac{F_{ph}}{c} = 72 \text{ cm}^{-3}, \quad (5)$$

where $d = 3.4$ kpc is the distance to Cas A. A similar reasoning gives a somewhat lower photon density for zone 2, but the photon field is much more anisotropic, on average coming from the direction of the remnant's center. For the spherical symmetry assumed, the anisotropy has no influence on the total inverse Compton emission, but as the maximum upscattering occurs for head on collisions, most energetic inverse Compton emission from zone 2 will be directed toward the center of the remnant, giving rise to a centrally brightened morphology, despite the fact that according to our model the dominant inverse Compton emission is coming from a narrow shell near the shock front.

In contrast, bremsstrahlung from the same electron population will show a ring-like morphology. Unfortunately, these interesting differences in morphology occur on the scale of arcminutes, which is currently beyond the reach of gamma-ray instruments.

Our two zone model is illustrated in Fig. 7, employing typical values for B and E_{break} near the shock front. The inverse Compton emission is completely dominated by the electrons in zone 2. In Fig. 7 inverse Compton emission is well below the TeV flux by *HEGRA*. It is therefore likely that the TeV emission is dominated by pion decay. However, we cannot yet make a definitive statement regarding the inverse Compton emission. For lower values of the magnetic field zone 2 is also the dominant source of bremsstrahlung by relativistic electrons. If the magnetic field is at the low end of our inferred magnetic field, $B = 0.08$ mG, and the far infrared photon density is larger than 150 cm^{-3} , the *HEGRA* detection, and the *BeppoSAX/CGRO-OSSE* upper limit on the non-thermal bremsstrahlung are still consistent with inverse Compton emission. The basic uncertainty in the photon density stems from the uncertain geometry, which for zone 2 may also include anisotropic photon emission. The combination of a low magnetic field and a rather high infrared photon density seems unlikely, but is not impossible. Even lower magnetic fields in zone 2 are unlikely, as the non-thermal bremsstrahlung would exceed the upper limit reported in §3.

7. CONCLUSION

We have provided estimates of the average magnetic field and the magnetic field near the shock front. The lower limit to the average magnetic field is based on the assumption that the electron spectrum follows a power law distribution in momentum, or is steeper than that at energies between 100-1000 keV.

The magnetic field strength near the shock front, was measured from the width of the narrow rim surrounding Cas A in X-ray continuum images, assuming that the emission mechanism is synchrotron emission, and the width is determined by synchrotron losses. The contribution of the X-ray synchrotron emission to the X-ray continuum flux is less than one third. This is supported by the X-ray morphology, which shows that even around 10 keV a substantial part of the continuum is associated with the bright ejecta shell (Bleeker et al. 2001).

This justifies taking into consideration alternative models than synchrotron radiation for the hard X-ray emission of Cas A. This was done in §2, where we tested the lower hybrid wave model. The bremsstrahlung spectrum predicted for electrons accelerated by lower hybrid waves has a similar spectral slope as is observed. The maximum energy to which the electrons are accelerated by this mechanism is ~ 95 keV, according to fits to the *BeppoSAX* data.

The estimated lower limit on the average magnetic field is 0.5 mG, whereas we find ~ 0.1 mG near the shock front. So even near the shock front the magnetic field strength is an order of magnitude above what is expected from shock compression of the canonical interstellar magnetic field of $3 \mu\text{G}$. This can be explained by either a higher magnetic

⁵ The other half of the photons escape directly, but all photons will have to pass through zone 2.

field strength in the medium surrounding Cas A, possibly as the result of the stellar wind of the progenitor (Biermann & Cassinelli 1993), or it means that the magnetic field is being rapidly enhanced near the shock front (Lucek & Bell 2000).

Electron energies close to the shock front may reach ~ 50 TeV or higher, but in contrast to Reynolds & Keohane (1999), we suggest that most of the cosmic ray electrons responsible for the bright radio ring are in an environment with a high magnetic field, where due to synchrotron losses the cut-off energy is well below the energy where X-ray synchrotron emission can be expected.

The high average magnetic field makes it likely that TeV emission from Cas A is dominated by pion decay (Atayan et al. 2000), but there is some uncertainty, as in our two zone model of Cas A it is still possible to obtain a high inverse Compton flux by assuming that the magnetic field is at the low end of what we have inferred from the X-ray rims, $B = 0.08$ mG, and using a higher far infrared photon density of 150 cm^{-3} .

Future mission will improve on those results. For instance, *GLAST* will probably be able to detect bremsstrahlung in the GeV range, which will provide a much more direct way of estimating the average magnetic field of Cas A.

As for the hard X-ray emission. *INTEGRAL* will not greatly improve the current measurement of the hard X-ray continuum emission, but by accurately measuring the ^{44}Ti related line emission, it will constrain that component

of the emission around 80 keV, and thus indirectly improve our understanding of the continuum emission around 100 keV.

One of the consequences of our analysis is that the gamma-ray emission will be mostly associated with the outer rim of Cas A, not with the bright X-ray and radio shell. Unfortunately, the spatial resolution required to measure this will not be available in the near future.

Finally, as an interesting by product of our analysis, we mention that the new modeling of the hard X-ray emission with the lower hybrid model yields revised values of the ^{44}Sc line emission at 68 keV and 78 keV, which correspond to a 50% higher mass of ^{44}Ti synthesized in the explosion compared to what was reported by Vink et al. (2001), i.e. $M(^{44}\text{Ti}) = 1.8 \times 10^{-4} M_{\odot}$. This is closer to the latest *CGRO-COMPTEL* measurements (Schönfelder et al. 2000).

JV was supported for this work by the NASA through Chandra Postdoctoral Fellowship Award Number PF0-10011 issued by the Chandra X-ray Observatory Center (CXO), which is operated by the Smithsonian Astrophysical Observatory (SAO) for and on behalf of NASA under contract NAS8-39073. JML was supported by basic research funds of the Office of Naval Research. We acknowledge the use of the HEASARC (Goddard Space Flight Center), CXO (SAO), and NCSA Astronomy Digital Image Library archival facilities.

APPENDIX

APPENDIX

In this appendix we want to demonstrate how free parameters in the fit of the model continuum (the maximum accelerated electron energy and the fraction of ambient electrons that are accelerated) are in principle determined by the plasma physics parameters of Cas A. The maximum electron energy is related to the maximum electric field that a lower hybrid wave can develop and the time that an electron can spend in this field, governed by the thickness of the shock precursor. The fraction of electrons that may be accelerated is expected to be determined by a marginal stability criterion, i.e. the electrons will be accelerated until the heating of the ambient plasma produced by Coulomb collisions of the accelerated electrons violates the conditions necessary for lower hybrid waves to exist, i.e. that the electron gyroradius be less than the wavelength divided by 2π .

The lower hybrid waves are generated by cross field ion motions produced by shocks. A fraction of preshock ions are reflected back upstream ahead of the shock with a velocity relative to the preshock plasma of $\sim 2v_s$, where v_s is the shock velocity. Ions reflected initially back along the shock velocity vector at a perpendicular shock travel a distance $0.68v_s/\Omega_i$ before returning (Gedalin 1996). Here Ω_i is the ion gyrofrequency. The maximum distance ahead of the shock that reflected ions can reach is $d = 1.37v_s/\Omega_i$. This obtains for ions reflected from the shock at an angle 30° to the shock front, so that $2v_s \sin 30^\circ = v_s$.

Consider a shock front at an angle θ to the local magnetic field direction. In order for the lower-hybrid waves generated ahead of the shock by the reflected ion precursor to grow to large amplitudes we require (Laming 2001b, i.e. the wave group velocity away from the shock is equal to the shock velocity, so that the time available for wave growth is essentially unlimited on the timescale of these processes)

$$\frac{\partial \omega}{\partial k_{\perp}} = \frac{v_s}{\cos \theta} = \frac{\alpha \omega}{k_{\perp}} \quad (\text{A1})$$

where $\alpha = 1/(2 \cos \beta \cos \theta)$ and β is the angle between $\partial \omega / \partial \vec{k}$ and the reflected ion bulk velocity vector $\vec{U}$. A minimum requirement for wave growth is $-1 < \alpha < -1/2$. Since it is ions *returning* to the shock that can generate the necessary waves, waves generated at the earliest possible period in the reflected ion orbit, i.e. lowest value of $|\cos \beta|$, when $\alpha \sim -1$, (for $\cos \theta \sim 1$) will dominate the growth.

The maximum electric field that the lower hybrid wave may develop before ion trapping and heating set is given by Karney (1978, in mks units);

$$E = B \left(\frac{\Omega_i}{\omega} \right)^{1/3} \frac{\omega}{4k_{\perp}} \simeq B \left(\frac{\Omega_i}{\omega} \right)^{1/3} \frac{v_s}{4 \cos \theta}. \quad (\text{A2})$$

Taking the wave frequency $\omega = \Omega_{LH} = \sqrt{\Omega_i \Omega_e}$, and the component of the electric field directed along the magnetic field direction to be $E\omega_{pi}/\omega_{pe}$ gives $a \simeq \Omega_{LH}v_s/16 \cos \theta$ for electron acceleration along the magnetic field direction. Ω_{LH} is the geometric mean of the ion and electron gyrofrequencies and is known as the lower hybrid frequency.

We treat the case of a nonrelativistic shock which may accelerate electrons to relativistic energies. We work in the rest frame of the preshock medium. In this frame an electron is accelerated by the component of the electric field directed along the magnetic field, until it outruns the reflected ion shock precursor a distance d from the shock front. Assuming acceleration begins with the electron at the shock front moving along $\vec{B}$ with initial velocity v_0 , it will continue for a time t given by

$$\left\{ \frac{c^2}{a} \sqrt{1 + \frac{(at + v_0/\sqrt{1 - v_0^2/c^2})^2}{c^2}} - \frac{c^2}{a\sqrt{1 - v_0^2/c^2}} \right\} \sin \theta = d + v_s t. \quad (A3)$$

which gives a quadratic equation for t :

$$t^2 (c^2 \sin^2 \theta - v_s^2) + t \left\{ \frac{2c^2}{a\sqrt{1 - v_0^2/c^2}} (v_0 \sin^2 \theta - v_s \sin \theta) - 2dv_s \right\} - d^2 - \frac{2dc^2 \sin \theta}{a\sqrt{1 - v_0^2/c^2}} = 0 \quad (A4)$$

We require $\sin \theta \geq v_s/v_0$ in order that the electron stays ahead of the shock front, and maximum acceleration time will be realized for $\sin \theta = v_s/v_0$. In this case $\cos \theta \simeq 1$ and the term in t in equation A4 simplifies to $-2dv_s t$ and

$$t = \frac{d}{v_s (c^2/v_0^2 - 1)} \left\{ 1 + \sqrt{1 + (c^2/v_0^2 - 1) \left(1 + \frac{2c^2}{adv_0 v_s \sqrt{1 - v_0^2/c^2}} \right)} \right\}. \quad (A5)$$

For $v_0 \ll c$, $at = \sqrt{2adv_0/v_s}$ and for $v_0 \sim c$, $at = 2ad/vs (c^2/v_0^2 - 1)$. Since $ad \propto v_s^2$ it is easy to see that the highest electron energies are most efficiently reached by fast shocks in hot plasma, i.e. for the highest plausible values of v_s and v_0 in Cas A. At $\sin \theta = v_s/v_0$, three to four shock interactions are required to accelerate electrons to the observed maximum energy of ~ 95 keV, increasing to about ten shock interactions if $\sin \theta = 1.2v_s/v_0$, for $v_s = 3000 \text{ km s}^{-1}$ and $v_0 = 3 \times 10^9 \text{ cm s}^{-1}$ (corresponding to an electron temperature of $4 \times 10^7 \text{ K}$). Even faster shock require fewer interactions. The maximum energy in the accelerated electron distribution will be determined by the number and obliquity of shock interactions experienced by these electrons during the evolution of Cas A, and so for the time being remains uncertain. However the maximum energy inferred from Fig. 1 is certainly plausible in these terms. In principle, given enough shock interactions, relativistic electrons can be produced. If a means of transporting these electrons to the forward shock were available, this might have a bearing on providing an injection mechanism for further electron acceleration by a Fermi mechanism.

The normalization of the non-thermal bremsstrahlung spectrum is determined by the fraction of plasma electrons that are accelerated in this way, and this will be determined ultimately by a marginal stability criterion. The instability requires the electron gyroradius to be less than the wavelength divided by 2π , and once a sufficient number of electrons are accelerated to heat the ambient plasma to temperatures higher than this, the instability will shut down. The ambient plasma loses energy through radiation, adiabatic expansion and possibly conduction. In Laming (2001b) the heating rate due to the observed accelerated electrons is estimated to be $4 \times 10^{-18} n_e n'_e \text{ ergs cm}^{-3} \text{ s}^{-1}$ for ambient and accelerated electron densities n_e and n'_e . The electron heating rate is $dT/dt = 0.019 n'_e \text{ K s}^{-1}$, and equating this to the cooling rate by adiabatic expansion of the Cas A remnant yields an accelerated electron fraction $n'_e/n_e \sim 0.02$. The inclusion of ionization and radiative power losses and relaxing the steady-state assumption inherent in this estimate will allow higher accelerated electron fractions.

In Laming (2001b) an accelerated electron fraction of 4% was found to give a good match to the ~ 50 ksec of BeppoSAX MECS and PDS data on Cas A then available. The estimate herein using only the ~ 500 ksec PDS data is higher at around 10% (see Fig. 1), but a range of plasma temperatures exists in Cas A (c.f. Vink et al. 1996; Willingale et al. 2001a), and only the hottest plasma with a temperature of $\sim 4 \times 10^7 \text{ K}$ contributes to the PDS bandpass. So with respect to all the hot plasma, an accelerated electron fraction of $\sim 4\%$ is consistent with the data, and preferred on theoretical grounds. Simulations of the evolution of the electron temperature following electron heating indicate that such an accelerated electron density heats plasma in the location of the contact discontinuity to temperatures in the range $4 - 5 \times 10^7 \text{ K}$. These temperatures are basically constant over the past ~ 30 years over which X-ray observations have been made. In this respect simulations of Cas A expanding into an $\rho \propto 1/r^2$ circumstellar medium appear to represent the physics much better than the simulations using a uniform density circumstellar medium in Laming (2001b).

REFERENCES

- | | |
|--|---|
| Aharonian, F., et al. 2001, <i>A&A</i> , 370, 112 | Atoyan, A., Tuffs, R. J., Aharonian, F. A., & Völk, H. J 2000, <i>A&A</i> , 354, 915 |
| Ahmad, I., et al. 1998, <i>PhRvL</i> , 80, 2550 | Arnaud, K.A. 1996, in <i>ASP Conf. Ser. 101, Astronomical Data Analysis Software and Systems V</i> ed. G. Jacoby & J. Barnes (San Francisco: ASP), 17 |
| Allen, G. E., et al. 1997, <i>ApJ</i> , 487, L97 | Ashworth, W.B 1980, <i>J. Hist. Ast.</i> , 11, 1 |
| Anders. E, & Grevesse, N. 1989, <i>Geochimica et Cosmochimica Acta</i> , 53, 197 | |
| Anderson, M.C., & Rudnick, L., 1996, <i>ApJ</i> , 456, 234 | |

- Asvarov, A.I., Dogiel, V.A., Guseinov, O.H., & Kasumov, F.K 1990, A&A, 229, 196
- Baars, J.W.M., Genzel, R., Pauliny-Toth, I. I. K., Witzel, A. 1977, A&A, 61, 99
- Balbus, S.A., & Hawley, J.F. 1991, ApJ, 400, 610
- Balbus, S.A., & Hawley, J.F. 1998, Rev. Mod. Phys., 70, 1
- Begelman, M.C., & Chiueh, T. 1988, ApJ, 332, 872
- Berezhko, E.G., Ksenofontov, L.T., & Voelk, H.J. 2002, preprint (astro-ph/0204085)
- Biermann, P.L., & Cassinelli, J.P. 1993, A&A, 691, 706
- Bingham, R., Kellett, B. J., Dawson, J. M., Shapiro, V. D., Mendis, D. A. 2000, ApJS, 127, 223
- Braun, R. 1987, A&A, 171, 233
- Bell, A.R. 1978, MNRAS, 182, 147
- Bleeker, J.A.M., Willingale, R., van der Heyden, K., Dennerl, K., Kaastra, J. S., Aschenbach, B., & Vink, J. 2001, A&A, 365, L225
- Blumenthal, G.R., & Gould, R.J. 1970, Rev. Mod. Physics, 42, 237
- Bykov, A.M., & Uvarov, Y.A. 1999, JETP, 88, 465
- Cowsik, R., & Sarkar, S. 1980, MNRAS, 191, 855
- Esposito, J.A., Hunter, D.A., Kanbach, G., & Skreekumar, P. 1996, ApJ, 461, 820
- Favata, F., et al. et al. 1997, A&A, 324, L49
- Fiore F., Guainazzi M., & Grandi P. 1999, "Handbook for BeppoSAX NFI Spectral Analysis", BeppoSAX Science Data Center, Rome
- Frontera, F., et al. 1997, A&AS, 122, 357
- Gedalin, M. 1996, JGR, 101, 4871
- Ginzburg, V.L., & Syrovatskii, S.I. 1965, ARA&A, 3, 297
- Görres, J., et al. 1998, PhRvL, 80, 2554
- Gotthelf, E.V., Koralesky, B., Rudnick, L., Jones, T. W., Hwang, U., & Petre, R. 2001, ApJ, 552, L39
- Gringauz, K. I., et al. 1986, Nature, 321, 282
- Haug, E. 1997, A&A, 326, 417
- Huba, J.D. "NRL Plasma formulary", 1994, Naval Research Laboratory, Washington DC
- Hughes, J., Rakowski, C., Burrows, D., & Slane, P. 2000, ApJ, 528, L109
- Hwang, U., Holt, S., & Petre, R. 2000, ApJ, 537, L119
- Iyudin A.F., et al. 1994, ApJ, 284, L1
- Johnson W.N., et al. 1993, ApJ, 693, 86
- Kaastra, J.S., Mewe, R., & Nieuwenhuizen, H. 1996, Proc. of the 11th Coll. on UV and X-ray, UV and X-ray Spectroscopy of Astrophysical and Laboratory Plasmas, ed. K. Yamashita & T. Watanabe (Tokyo:Universal Academy Press), 411
- Karney, C. F. F. 1978, Phys. Fluids, 21, 1584
- Klimov, S. S., et al. 1986, Nature, 321, 292
- Koyama, K., Petre, R., Gotthelf, E. V., Hwang, U., Matsuura, M., Ozaki, M., & Holt, S. S. 1995, Nature, 378, 255
- Krasnosel'skikh, V. V., Kruchina, E. N., Thejappa, G., & Volokitin, A. S. 1985, A&A, 149, 323
- Lagage, P. O., Claret, A., Ballet, J., Boulanger, F., Cesarsky, C. J., Cesarsky, D., Fransson, C., & Pollock, A. 1996, A&A, 315, L273
- Laming, J.M., 2001a ApJ, 546, 1149
- Laming, J.M., 2001b ApJ, 563, 828
- Longair, M.S., High Energy Astrophysics, Vol. 2: Stars, the galaxy and the interstellar medium 1994 (2nd ed.; Cambridge:Cambridge University Press)
- Luceck, S.G., & Bell, A.R. 2000, ApJ, 314, 65
- Norman, E.B., et al. 1998, PhRvC, 57, 2010
- Nozawa, S., Itoh, N., & Kohyama, Y. 1998, ApJ, 507, 530
- Quataert, E., Dorlan, W., & Hammet, G. 2002, ApJ, in press, preprint astro-ph/0205492()
- Reed, J.E., Hester, J.J., Fabian, A.C., & Winkler, P.F. 1995, ApJ, 440, 706
- Reynolds, S.P. 1998, ApJ493, 375
- Reynolds, S.P., & Keohane, J.W. 1999, ApJ, 525, 368
- Schönfelder, V., et al. 2000, in AIP Conf. Proc. 510, The Fifth Compton Symposium, ed. M.L. McConnell & J.M. Ryan (Melville, NY: AIP), 54
- Shapiro, V. D., Bingham, R., Dawson, J. M., Dobe, Z., Kellett, B. J., & Mendis, D. A. 1999, J. Geophys. Res., 104, 2537
- Strong, A. W., Bloemen, H., Collmar, W., Diehl, R., Hermesen, W., Iyudin, A., Schönfelder, V., & The, L.-S. 2000, in AIP Conf. Proc. 510, The Fifth Compton Symposium, ed. M.L. McConnell & J.M. Ryan (Melville, NY: AIP), 60
- Tanimori, T., et al. 1998, ApJ, 497, L25
- The, L.-S., Leising, M.D., Kurfess, J.D., Johnson, W. N., Hartmann, D.H., Gehrels, N., Grove, J. E., & Purcell, W. R. 1996, A&AS, 120, 357
- Thompson A., et al., (editors), "X-ray data booklet" 2001, Lawrence Berkeley National Laboratory
- Vaisberg, D. L., Galeev, A. A., Zastenker, G. N., Klimov, S. I., Nozdachev, M. N., Sagdeev, R. Z., Sokolov, A. Y., & Shapiro, V. D., 1983, Sov. Phys. JETP, 58, 716
- Vink, J., Kaastra, J.S., & Bleeker, J.A.M. 1996, A&A, 307, L41
- Vink, J., Bloemen, H., Kaastra, J.S., & Bleeker, J.A.M. 1998, A&A, 339, 201
- Vink, J., Laming, J. M., Kaastra, J.S., Bleeker, J. A. M., Bloemen, H., & Oberlack, U. 2001, ApJ, 560, L79
- Vink, J., & Laming, J.M. 2002, Proc. of the XXII Moriond Astrophysics Meeting, The Gamma-Ray Universe, ed. A. Goldwurm, D. Neumann, & J. Tran Thanh Van (astro-ph/0205235)
- Willingale, R., Bleeker, J.A.M., van der Heyden, K.J., Kaastra, J. S., & Vink, J. 2002, A&A, 381, 1039
- Willingale, R., Bleeker, J.A.M., van der Heyden, K.J., & Kaastra, J. S. 2001b, A&A, submitted, preprint(astro-ph/0207273)
- Wright, M., Dickel, J., Koralesky, B., & Rudnick, L. 1999, ApJ, 518, 284

TABLE A1
SUMMARY OF MODEL FITS TO THE *BeppoSAX*-PDS SPECTRUM.

	LHW model	<i>srcut</i>	<i>sresc</i>	power law
Norm continuum ^a	0.058 ± 0.002	2522	2522	2.37 ± 0.2
power law index				-3.32 ± 0.05
E_{max}/E_{cutoff} (keV)	95 ± 5	1.21 ± 0.01	3.42 ± 0.02	
Norm thermal emission ^b	0.62 ± 0.02	7.6 ± 3.3	0.0(< 2.5)	0.0(< 17)
⁴⁴ Sc line flux (10 ⁻⁵ phcm ⁻² s ⁻¹)	3.2 ± 0.3	2.7 ± 0.4	2.9 ± 0.4	2.1 ± 0.4
Ta K α flux (10 ⁻⁵ phcm ⁻² s ⁻¹)	2.4 ± 0.7	2.6 ± 0.8	2.8 ± 0.8	1.4 ± 0.
$\chi^2/d.o.f$	273.1/251	273.0/252	289.3/252	267.5/251

^a The normalization for the LWH model is defined as $10^{12} n_H n_e V / 4\pi d^2 \text{ cm}^{-5}$. The normalization for the synchrotron models is fixed to the radio flux density at 1 GHz for the epoch 2000.0, and radio spectral index is 0.775. These values are based on a compilation of radio data, and includes the effect of the secular flux decrease of Cas A (Baars et al. 1977). The normalization for the power law model is ph/s/keV at 1 keV.

^b The normalization is defined as $10^{12} n_H n_e V / 4\pi d^2$. The temperature was fixed to 4×10^7 K. Note that this definition assumes solar abundances, which is not applicable for Cas A, so care should be taken before using it to calculate electron densities.

TABLE A2
BEST FIT PARAMETERS FOR SPECTRA FROM THE NORTHEASTERN RIM.

	NE Rim	Inside NE Rim
kT (keV)	3.7 ± 0.7	3.0 ± 0.4
$n_e t$ ($10^{10} \text{ cm}^{-3} \text{ s}^{-1}$)	3.3 ± 0.5^a	5.2 ± 0.6
Ne		$0.4^{+1.0}_{-0.4}$
Mg		0.9 ± 0.4
Si		4.5 ± 0.6
S		4.3 ± 0.7
Ar		4.4 ± 1.5
Ca		5.4 ± 2.5
Fe		2.2 ± 0.4
EM c (10^{10} cm^{-5})	3.0 ± 0.4	7.5 ± 1.0
PL index	-2.24 ± 0.05	$-4.5 (< -4.3)^d$
PL norm ($10^{-3} \text{ s}^{-1} \text{ cm}^{-2} \text{ }^e$)	2.5 ± 0.2	3.4 ± 0.6
PL contribution 4 - 6 keV	84%	11%
N_H (10^{22} cm^{-2})		1.08 ± 0.03
χ^2/ν	261/161	363/153

Note. — The spectral model consisted of *SPEX*'s non-equilibrium ionization model (v1.10), and a power law (PL) continuum, in order to approximate the synchrotron radiation. The column density and abundances were first determined for the spectrum inside the NE Rim, and subsequently applied to the rim spectrum. Following Vink et al. (1996) we set He and N abundances to 10 times solar (Anders & Grevesse 1989). The poorly constrained O abundance was coupled to Si; Ni was coupled to Fe. Errors correspond to 95% confidence intervals.

^aNote that this value is roughly consistent with the synchrotron loss times, if the electron density is $n_e \sim 30 \text{ cm}^{-3}$.

^bThe He and N abundances were assumed to be 10 times solar, following Vink et al. (1996).

^cThe emission measure (EM) is defined as $n_H n_e V / 4\pi d^2$.

^dThe power law slope was poorly constrained, but tended to be steep, we therefore fixed it to a value of 4.5. The lower limit corresponds to $\Delta\chi^2 = 4$.

^eAt 1 keV.